

# Potential for using climate forecasts in spatio-temporal prediction of dengue fever incidence in Malaysia

Submitted by

**Norziha Che Him**

to the University of Exeter as a thesis for the degree of Doctor of Philosophy in Mathematics, October 2015.

This thesis is available for Library use on the understanding that it is copyright material and that no quotation from the thesis may be published without proper acknowledgement.

I certify that all material in this thesis which is not my own work has been identified and that no material is included for which a degree has previously been conferred upon me.

.....  
Norziha Che Him

# Abstract

Dengue fever is a viral infection transmitted by the bite of female *Aedes aegypti* mosquitoes. It is estimated that nearly 40% of the world's population is now at risk from Dengue in over 100 endemic countries including Malaysia. Several studies in various countries in recent years have identified statistically significant links between Dengue incidence and climatic factors. There has been relatively little work on this issue in Malaysia, particularly on a national scale. This study attempts to fill that gap. The primary research question is 'to what extent can climate variables be used to assist predictions of dengue fever incidence in Malaysia?'. The study proposes a potential framework of modelling spatio-temporal variation in dengue risk on a national scale in Malaysia using both climate and non-climate information.

Early chapters set the scene by discussing Malaysia and Climate in Malaysia and reviewing previous work on dengue fever and dengue fever in Malaysia. Subsequent chapters focus on the analysis and modelling of annual dengue incidence rate (DIR) for the twelve states of Peninsular Malaysia for the period 1991 to 2009 and monthly DIR for the same states in the period 2001 to 2009.

Exploratory analyses are presented which suggest possible relationships between annual and monthly DIR and climate and other factors. The variables that were considered included annual trend, in year seasonal effects, population, population density and lagged dengue incidence rate as well as climate factors such as average rainfall and temperature, number of rainy days, ENSO and lagged values of these

climate variables. Findings include evidence of an increasing annual trend in DIR in all states of Malaysia and a strong in-year seasonal cycle in DIR with possible differences in this cycle in different geographical regions of Malaysia. High population density is found to be positively related to monthly DIR as is the DIR in the immediately preceding months. Relationships between monthly DIR and climate variables are generally quite weak, nevertheless some relationships may be able to be usefully incorporated into predictive models. These include average temperature and rainfall, number of rainy days and ENSO. However lagged values of these variables need to be considered for up to 6 months in the case of ENSO and from 1-3 months in the case of other variables.

These exploratory findings are then more formally investigated using a framework where dengue counts are modelled using a negative binomial generalised linear model (GLM) with a population offset. This is subsequently extended to a negative binomial generalised additive model (GAM) which is able to deal more flexibly with non-linear relationships between the response and certain of the explanatory variables. The model successfully accounts for the large amount of overdispersion found in the observed dengue counts. Results indicated that there are statistically significant relationships with both climate and non-climate covariates using this modelling framework. More specifically, smooth functions of year and month differentiated by geographical areas of the country are significant in the model to allow for seasonality and annual trend. Other significant covariates included were mean rainfall at lag zero month and lag 3 months, mean temperature at lag zero month and lag 1 month, number of rainy days at lag zero month and lag 3 months, sea surface temperature at lag 6 months, interaction between mean temperature at lag 1 month and sea surface temperature at lag 6 months, dengue incidence rate at lag 3 months and population density.

Three final competing models were selected as potential candidates upon which an early warning system for dengue in Malaysia might be able to be developed. The model fits for the whole data set were compared using simulation experiments to allow for both parameter and negative binomial model uncertainty and a single

model preferred from the three models was identified. The ‘out of sample’ predictive performance of this model was then compared and contrasted for different lead times by fitting the model to the first 7 years of the 9 years monthly data set covering 2001-2009 and then analysing predictions for the subsequent 2 years for lead time of 3, 6 12 and 24 months. Again simulation experiments were conducted to allow for both parameter and model uncertainty. Results were mixed. There does seem to be predictive potential for lead times of up to six months from the model in areas outside of the highly urbanised South Western states of Kuala Lumpur and Selangor and such a model may therefore possibly be useful as a basis for developing early warning systems for those areas. However, none of the models developed work well for Kuala Lumpur and Selangor where there are clearly more complex localised influences involved which need further study.

This study is one of the first to look at potential climatic influences on dengue incidence on a nationwide scale in Malaysia. It is also one of the few studies worldwide to explore the use of generalised additive models in the spatio-temporal modelling of dengue incidence. Although, the results of the study show a mixed picture, hopefully the framework developed will be able to be used as a starting point to investigate further if climate information can valuably be incorporated in an early warning system for dengue in Malaysia.



# Acknowledgements

My thanks to my supervisor Professor Trevor Bailey for his support and guidance through my PhD journey in CEMPS at the University of Exeter. Thanks also to Professor David Stephenson and Dr Christopher Ferro as, respectively, my second supervisor and postgraduate mentor in CEMPS. Special thanks to all Exeter Climate System academics and administrative support staff who helped me in various ways including Dr Catherine, Dr Rachel, Dr Alemtsehai, Dr Theo, Dr Stan and Dr Yuwei. Lastly, to all friends - Dr Ummu Atiqah, Dr Asma, Dr Alasdair, Dr Lester, Dr Maria, Dr Robin, Siti Hasmah and Noraliza.

Thank you to the Ministry of Education Malaysia (MOE) and Universiti Tun Hussein Onn Malaysia (UTHM) for my financial support in respect of this study. Appreciation to Dr Lokman Hakim, Director of Disease Control Division, for his kind assistance regarding dengue data, also to the Malaysian Meteorological Department (MMD) and the Department of Irrigation and Drainage Malaysia (DIDM) for supplying temperature and rainfall data.

Special thanks to my husband, Yusliandy Yusof, my adorable sons and daughter; Afham Hakim, Ammar Hafiz and Amna Haura for their love and support. Appreciation also to my mother Siti Melia Awang and my late father, Che Him Mamat, for their encouragement in my studies. Lastly, thanks to my family in law and all my brothers and sisters, Mazrah, Zaidi, Al-Zuhari, Azani, Azman, Ariza, Nor-Roshayati, Kamarul Halim, Khairiah Mahayu, Kamarudin and Kamariah, thank you all for your love and your invaluable encouragement.

# Contents

<b>Acknowledgements</b>	<b>5</b>
<b>Contents</b>	<b>6</b>
<b>List of Figures</b>	<b>9</b>
<b>List of Tables</b>	<b>16</b>
<b>1 Introduction</b>	<b>18</b>
1.1 Motivation . . . . .	18
1.2 Research Aims . . . . .	25
1.3 Summary . . . . .	27
<b>2 Malaysia and its climate</b>	<b>29</b>
2.1 Introduction to Malaysia . . . . .	29
2.1.1 Demographic, Socio-economic and Health Profile . . . . .	32
2.1.2 Sources of Cartographic and Demographic data . . . . .	36
2.2 Climate in Malaysia . . . . .	38

2.2.1	General Description of Climate . . . . .	39
2.2.2	Sources of climate data . . . . .	42
2.3	Summary . . . . .	44
<b>3</b>	<b>Dengue Fever and Dengue Fever in Malaysia</b>	<b>45</b>
3.1	Dengue Fever and its transmission . . . . .	45
3.2	Dengue Fever worldwide . . . . .	49
3.3	Dengue Fever in Malaysia . . . . .	56
3.3.1	Distribution of dengue in Malaysia . . . . .	57
3.3.2	Surveillance and control . . . . .	62
3.3.3	Sources of dengue data . . . . .	66
3.4	Climate and dengue . . . . .	67
3.4.1	Global perspective . . . . .	68
3.4.2	Malaysia perspective . . . . .	74
3.5	Summary . . . . .	76
<b>4</b>	<b>Exploratory data analysis</b>	<b>77</b>
4.1	Description of collated datasets . . . . .	77
4.2	Patterns in Dengue Incidence . . . . .	80
4.2.1	Annual Patterns in Dengue Incidence . . . . .	83
4.2.2	Monthly Patterns in Dengue Incidence . . . . .	93
4.3	Dengue Incidence and Demographic Data . . . . .	102

4.4	Dengue Incidence and Climate Data . . . . .	104
4.5	Summary . . . . .	118
<b>5</b>	<b>Model Development</b>	<b>121</b>
5.1	Modelling Frameworks . . . . .	121
5.2	Covariate Selection . . . . .	131
5.3	Model Comparison and Development . . . . .	134
5.4	Summary . . . . .	151
<b>6</b>	<b>Model Testing</b>	<b>154</b>
6.1	Model Testing — Comparison of overall fit . . . . .	156
6.2	Model Testing — Out of sample predictions . . . . .	165
6.3	Model Testing — Out of sample predictions (Kuala Lumpur/Selangor)	168
6.4	Summary . . . . .	176
<b>7</b>	<b>Summary and Conclusions</b>	<b>177</b>
7.1	Main findings . . . . .	177
7.2	Limitations of the study . . . . .	181
7.3	Scope for future work . . . . .	182
7.4	Summary . . . . .	183
	<b>References</b>	<b>185</b>



# List of Figures

2.1	<i>Map of Peninsular Malaysia (left) and East Malaysia (right).</i>	31
2.2	<i>Map of the 12 states in the Peninsular of Malaysia.</i>	36
2.3	<i>Map showing five regions (referred to as Niño 1, Niño 2, Niño 3, Niño 3.4 and Niño 4) in the Pacific identified as important locations for monitoring winds, sea surface temperatures and rainfall activities, changes that may be associated to varying degrees with El Niño process.</i>	42
3.1	<i>Countries at risk of dengue in 2010.</i>	50
3.2	<i>(a) The total number of dengue cases, (b) The total population size and (c) DIR per 100,00 population in Malaysia from 1991 to 2014 (Source : Ministry of Health Malaysia, 2014).</i>	59
4.1	<i>Annual population estimates for the four regions of Malaysia from 2001 to 2009.</i>	82
4.2	<i>Annual DIR per 100,000 population for the 12 states of Peninsular Malaysia from 1991-2009.</i>	84
4.3	<i>Annual DIR per 100,000 population for the four regions (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 1991-2009.</i>	85

4.4	<i>Annual DIR per 100,000 population for main regions of Malaysia considered in this study.</i>	86
4.5	<i>Annual DIR per 100,000 population for 3 states (a) Kelantan, (b) Terengganu and (c) Pahang in the North East region from 1991 to 2009 for Malaysia.</i>	88
4.6	<i>Annual DIR per 100,000 population for 3 states (a) Perlis, (b) Penang and (c) Kedah in the North West region from 1991 to 2009 for Malaysia.</i>	89
4.7	<i>Annual DIR per 100,000 population for 3 states (a) Negeri Sembilan, (b) Melaka and (c) Johor in the South East region from 1991 to 2009 for Malaysia.</i>	90
4.8	<i>Annual DIR per 100,000 population for 3 states (a) Perak, (b) Selangor and (c) Kuala Lumpur in the South West region from 1991 to 2009 for Malaysia.</i>	91
4.9	<i>Average monthly DIR values for Malaysia 2001-2009.</i>	93
4.10	<i>Monthly DIR per 100,000 population for North East, South East, North West and South West of Malaysia from 2001 to 2009.</i>	95
4.11	<i>The mean annual cycle of monthly DIR per 100,000 population for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009.</i>	96
4.12	<i>The mean monthly cycle of DIR per 100,000 population for 3 states; (a) Kelantan, (b) Terengganu and (c) Pahang in the North East region in Malaysia from 2001 to 2009.</i>	98
4.13	<i>The mean monthly cycle of DIR per 100,000 population for 3 states; (a) Perlis, (b) Penang and (c) Kedah in the North West region in Malaysia from 2001 to 2009.</i>	99

4.14	<i>The mean monthly cycle of DIR per 100,000 population for 3 states; (a) N.Sembilan, (b) Melaka and (c) Johor in the South East region in Malaysia from 2001 to 2009. . . . .</i>	100
4.15	<i>The mean monthly cycle of DIR per 100,000 population for 3 states; (a) Perak, (b) Selangor and (c) Kuala Lumpur in the South West region in Malaysia from 2001 to 2009. . . . .</i>	101
4.16	<i>Relationship between log monthly DIR and population density for 12 states of Malaysia 2001-2009 . . . . .</i>	103
4.17	<i>Relationship between log monthly DIR for 12 states of Malaysia 2001-2009 and (a) log monthly DIR lag 1 Month, (b) log monthly DIR lag 2 Months, (c) log monthly DIR lag 3 Months. . . . .</i>	105
4.18	<i>Relationship between log monthly DIR for 12 states of Malaysia 2001-2009 and (a) Rainfall Current Month, (b) Rainfall Lag 1 Month, (c) Rainfall Lag 2 Months and (d) Rainfall Lag 3 Months. . . . .</i>	106
4.19	<i>Relationship between log monthly DIR for 12 states of Malaysia 2001-2009 and (a) Number of Rainy Days Current Month, (b) Number of Rainy Days Lag 1 Month, (c) Number of Rainy Days Lag 2 Months and (d) Number of Rainy Days Lag 3 Months. . . . .</i>	107
4.20	<i>Relationship between log monthly DIR for 12 states of Malaysia 2001-2009 and (a) Temperature Current Month, (b) Temperature Lag 1 Month, (c) Temperature Lag 2 Months and (d) Temperature Lag 3 Months. . . . .</i>	108
4.21	<i>Relationship between log monthly DIR for 12 states of Malaysia 2001-2009 and (a) Humidity Current Month, (b) Humidity Lag 1 Month, (c) Humidity Lag 2 Months and (d) Humidity Lag 3 Months. . . . .</i>	109
4.22	<i>Relationship between log monthly DIR and average rainfall in same month for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	110

4.23	<i>Relationship between log monthly DIR and average rainfall 3 months previously for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	111
4.24	<i>Relationship between log monthly DIR and number of rainy days in same month for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	112
4.25	<i>Relationship between log monthly DIR and number of rainy days 3 months previously for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	113
4.26	<i>Relationship between log monthly DIR and average temperature in same month for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	114
4.27	<i>Relationship between log monthly DIR and average temperature 3 months previously for (a) North East, (b) South East, (c) North West and (d) South West of Malaysia from 2001 to 2009. . . . .</i>	115
4.28	<i>Mean annual cycle of (a) log monthly DIR, (b) Average Rainfall, (c) Average Temperature (d) No. of Rainy Days in South West of Malaysia from 2001 to 2009. . . . .</i>	116
4.29	<i>Standardised anomalies of SST (Niño 4 Index) (3 months-running mean) (Jan 2001-Dec 2009). . . . .</i>	118
4.30	<i>Relationship between log monthly DIR and Niño 4 at different lags in the 12 states of Peninsular Malaysia from 2001 to 2009. . . . .</i>	119
5.1	<i>(a) Residuals vs. fitted plot, (b) Absolute value of Residuals vs. fitted plot, (c) Residuals vs. Theoretical Quantiles plot and (d) Frequency vs. Residuals plot. . . . .</i>	138

- 5.2 Smooth annual cycle functions of four regions in Malaysia; upper row from left to right are North East and North West region, and lower row from left to right are South East and South West region. In each panel, the solid line is the estimate, and the dashed line is the confidence interval. 139
- 5.3 Smooth functions in Model A. Upper row from left to right are North East, North West, South East and South West region for smooth function of month (by region). Lower row from left to right are North East, North West, South East and South West region for smooth function of year (by region). In each panel, the solid line is the estimate, and the dashed line is the confidence interval. . . . . 145
- 5.4 Smooth functions in Model B. Upper row from left to right are monsoon Southwest and Northeast for smooth function of month (by monsoon) followed by North East for smooth function of year (by region). Lower row from left to right are North West, South East and Sourelnship with th West for smooth function of year (by region). In each panel, the solid line is the estimate, and the dashed line is the confidence interval. . 146
- 5.5 Smooth functions in Model C. First three rows are smooth functions of month for each of the twelve states. Last row is smooth function of year for the four regions. In each panel, the solid line is the estimate, and the dashed line is the confidence interval. . . . . 147
- 5.6 Comparison of most significant coefficients (a) Rainfall and Rainfall Lag 3, (b) Rainy Days, Rainy Day Lag 3 and DIR Lag 3, (c) Temperature and (d) Population Density for Model A, Model B and Model C with estimates error bars represent 95% confidence intervals around the respective mean values. . . . . 152
- 6.1 Monthly DIR fitted values for 2001-2009 for Peninsular Malaysia for Models A, B and C and associated root mean square error (RMSE) of constituent fitted values for each month and for each state. . . . . 158

6.2	<i>Q-Q plot of standardised deviance residuals for Model A for each state and month from 2001 to 2009. . . . .</i>	158
6.3	<i>Q-Q plot of standardised deviance residuals for Model B for each state and month from 2001 to 2009. . . . .</i>	159
6.4	<i>Q-Q plot of standardised deviance residuals for Model C for each state and month from 2001 to 2009. . . . .</i>	159
6.5	<i>Fitted versus observed values of monthly DIR from Model A for each state and each month for 2001-2009 and associated RMSE. . . . .</i>	160
6.6	<i>Monthly DIR fitted values for 2001-2009 for Peninsular Malaysia for Model A and associated simulated 95% CIs for mean values for each month and for each state. . . . .</i>	162
6.7	<i>Monthly DIR fitted values for 2001-2009 for Peninsular Malaysia for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .</i>	163
6.8	<i>Fitted versus observed values of monthly DIR from Model A for each state and each month for North East region for 2001-2009 and associated RMSE.</i>	163
6.9	<i>Fitted versus observed values of monthly DIR from Model A for each state and each month for South West region for 2001-2009 and associated RMSE.</i>	164
6.10	<i>Monthly DIR fitted values for 2001-2009 for South West region for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .</i>	164
6.11	<i>Monthly DIR out of sample predicted values for 2008-2009 for Peninsular Malaysia for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .</i>	166

- 6.12 *Out of sample predicted values for 2008-2009 for each month and for each state versus observed values for Peninsular Malaysia for Model A and associated RSME values for lead times of 3, 6, 12, and 24 months. . . . .* 167
- 6.13 *Monthly DIR out of sample predicted values for 2008-2009 for North East region for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .* 168
- 6.14 *Out of sample predicted values for 2008-2009 for each month and for each state versus observed values for North East Malaysia for Model A and associated RSME values for lead times of 3, 6, 12, and 24 months. . . . .* 169
- 6.15 *Monthly DIR out of sample predicted values for 2008-2009 for South West region for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .* 170
- 6.16 *Out of sample predicted values for 2008-2009 for each month and for each state versus observed values for South West Malaysia for Model A and associated RSME values for lead times of 3, 6, 12, and 24 months. . . . .* 171
- 6.17 *Monthly DIR out of sample predicted values for 2008-2009 for Peninsular Malaysia without the states of Kuala Lumpur and Selangor for Model A and associated simulated 95% prediction intervals for each month and for each state. . . . .* 172
- 6.18 *Out of sample predicted values for 2008-2009 for each month and for each state versus observed values for Peninsular Malaysia without the states of Kuala Lumpur and Selangor for Model A and associated RSME values for lead times of 3, 6, 12, and 24 months. . . . .* 173
- 6.19 *Out of sample predicted values for 2008-2009 for each month and for each state versus observed values for Peninsular Malaysia without the states of Kuala Lumpur and Selangor for Model A and associated RSME values for lead times of 3, 6, 12, and 24 months. . . . .* 174



# List of Tables

2.1	<i>Malaysia top 10 causes of death.</i>	35
2.2	<i>Distribution of area, latitude, longitude and population in Malaysia.</i>	37
4.1	<i>The division of the states in Malaysia.</i>	79
4.2	<i>Source and original resolution of dataset.</i>	81
4.3	<i>Total number of dengue cases and average DIR in each region from 1991-2009 for Malaysia.</i>	87
4.4	<i>State with highest annual DIR from 2001-2009.</i>	92
4.5	<i>Notified total of dengue and maximum monthly DIR of main regions in Malaysia from 2001-2009.</i>	95
5.1	<i>Final Selected Variables from negative binomial GLM runs.</i>	135
5.2	<i>Likelihood statistic (<math>L</math>), degrees of freedom (<math>n - p</math>), Deviance (<math>D</math>), Akaike Information Criterion (AIC) and Bayesian Information Criterion (BIC) for models with similar subsets of covariates fitted using Poisson and negative binomial GLMs.</i>	136
5.3	<i>Estimates of coefficients, standard errors and <math>p</math>-values of negative binomial GLM model.</i>	137



5.4	<i>Comparing the negative binomial GLM and GAM: ANOVA results. . . .</i>	139
5.5	<i>Deviance, pseudo-<math>R_D^2</math>, number of parameter(<math>p</math>), degrees of freedom (<math>n-p</math>), AIC and BIC for models with different subsets of covariates fitted using the negative binomial GAM. . . . .</i>	141
5.6	<i>Parameter estimates, standard errors and p-values for negative binomial GAM Model 5.28. . . . .</i>	142
5.7	<i>The deviance (<math>D</math>), log-likelihood, Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC) and Un-Biased Risk Estimator (UBRE) for Model A, Model B and Model C conditioned by region, monsoon and state respectively using negative binomial GAM. . . . .</i>	144
5.8	<i>Estimates of the dengue count Model A (by region) parameters. . . . .</i>	148
5.9	<i>Estimates of the dengue count Model B (by monsoon) parameters. . . . .</i>	149
5.10	<i>Estimates of the dengue count Model C (by state) parameters. . . . .</i>	150
6.1	<i>Log-likelihood (LogLik), Akaike Information Criterion (AIC) and effective degrees of freedom (EDF) for Model A, Model B and Model C for overall fit to the full 108 month data set . . . . .</i>	157

# Chapter 1

## Introduction

This chapter introduces the key motivations behind this study and the primary research aims which are addressed. The focus of the study is on exploring the relationship between climatic variables and the incidence of dengue fever in Malaysia and also the potential for using any such relationships to assist in providing early warning forecasts of dengue epidemics. The chapter starts by outlining the growing need to better understand the risk factors associated with dengue fever, both globally and, more specifically, in the context of Malaysia. It then goes on to specify research aims for the remainder of the thesis and concludes by setting out a structure for the subsequent chapters of the study.

### 1.1 Motivation

Dengue fever (DF) is a viral infection characterised by sudden high fever, severe headache, rash, muscle and joint pains. The virus is transmitted by the bite of female *Aedes aegypti* mosquitoes and infection rates of dengue can be as high as 90% among those who have not been previously exposed to the virus (Gubler, 1998). Dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS) are more serious (and potentially fatal) complications of the disease. Guzman and Kouri

(2002) have reported that nearly 40% of the world's population is now estimated to be at risk from DF in over 100 endemic countries and 500,000 people are estimated to be hospitalised every year with DHF.

The number of dengue cases in Malaysia continues to rise annually and DF is now recognised as a significant public health problem in that country (Smith, 1957; Aziz et al., 2012; Chew et al., 2012). Efforts to reduce the number of dengue cases is now a high priority of various internal and external agencies in Malaysia, not least the Malaysian Ministry of Health which has the main responsibility in addressing the situation. Dengue fever was first reported in Malaysia by Skae (1902), followed by dengue hemorrhagic fever and dengue shock syndrome epidemics in 1962 in Penang, and dengue fever cases first became officially notifiable in 1971 (Rudnick et al., 1965; Poovaneswari, 1993). Now in Malaysia it is the responsibility of all medical practitioners to report every case of dengue fever to the nearest Local Health Office within 24 hours from the time it was diagnosed (Narwani et al., 2005). As the reporting systems have developed, and particularly since 1980, the Malaysian Ministry of Health has recorded continual rising annual cases of dengue to the extent that Ang et al. (2010) recently highlighted dengue fever as an urgent major public health threat in the highly urbanised states of Selangor and Kuala Lumpur.

In general, climate is known to have the potential to influence human health through both direct and indirect mechanisms. The direct mechanisms include, for example, episodes of heat or cold stress and extreme events (drought and flood), while the indirect mechanisms include, for example, the impact of climate anomalies on the risk of vector borne infectious diseases such as malaria and dengue through changing environmental conditions for the vector (Connor et al., 2010). According to Gage et al. (2008), prediction of the relative impact of sustained climate change for vector borne diseases is difficult and will require long-term studies that need to look not only at the effects of climate change but also the contributions of other agents of global change. That said, several studies worldwide have revealed relationships between climatic variables and dengue fever and how these interact

with other known risk factors such as socio-economic conditions. Such studies have typically used statistical modelling methods of varying sophistication including time-series analysis, multivariate regression, generalised linear models (GLM), generalised linear mixed models (GLMM), generalised additive models (GAM) or generalised additive mixed models (GAMM). Better understanding of how socio-economic and climatic factors can affect the transmission of dengue fever may help in developing early warning systems (EWS) for epidemics of dengue and so widen the effectiveness of responsive measures (surveillance and prevention). Khun et al. (2005) emphasised the importance of developing systems for early identification for dengue epidemics to help health authorities in surveillance and prevention. To be effective such EWS need to be able to target forecasts geographically within a country into smaller areas such as districts, states or regions (Hu et al., 2012).

In Malaysia, the dengue incidence rate (DIR) in the country as a whole in recent years has fluctuated from as low as 27.5 cases per 100,000 population in 1995 to the high level of 132.5 cases per 100,000 population in 2004 (Kumarasamy, 2006). Epidemics of dengue have occurred roughly every four years with major outbreaks recorded in 1974, 1978, 1982 and 1990 (Lam, 1993b) and with a generally similar pattern with increasing incidence since then. Significant work has been introduced since the 1970s on prevention and control programmes to eliminate the *Aedes* mosquitoes and larval breeding habitats and on public education and law enforcement. However, there have been relatively few modelling studies on the relationship between dengue and climatic and other risk factors in Malaysia as a whole and even less work on the practical development of EWS. That background provides the primary motivation for the work developed throughout the subsequent chapters of this study.

One key challenge in pursuing that agenda is the availability of limited information at the local geographical scale in Malaysia, for example models considered in Racloz et al. (2012) were unable to sufficiently account for the spatio-temporal features of the disease because of the limited geographical resolution of available covariates. However, although there has been little work on dengue EWS in Malaysia, there

has been more progress in the neighbour of Malaysia to the south. Dengue fever in Singapore was first recognised as an important public health issue in the early 1960s and *Aedes* control programs have been in place since 1969. Forecasting models of dengue fever have also been developed in Singapore. Ma et al. (2008) carried out a study to look at the association between socio-economic variables and dengue incidence in Singapore for five years from 1998 to 2002 and identified significant association between dengue cases and socio-economic or demographic variables, with areas of higher proportion of disadvantaged residents having more dengue cases. Another more recent study in Singapore used Poisson time series modelling including climate factors such as rainfall and temperature up to 16 weeks or 4 months in advance (Hii et al., 2012). Such studies could be the benchmark for encouraging further research, such as that intended in this study, into the relationship between dengue and climatic variables and other risk factors in Malaysia and the potential for developing EWS for dengue based upon such relationships.

Having established that basic motivation, it is perhaps useful at this point to provide more specific detail on some of the issues so far raised; firstly, in relation to dengue on the world stage and, secondly, in the specific context of Malaysia.

As said previously, dengue fever is a vector borne viral infection transmitted by the bite of female *Aedes aegypti* mosquitoes. The number of dengue cases has increased dramatically around the world in recent years due to the absence of vaccines and drugs (WHO, 2012b). The illness is caused by one of four strains of the dengue virus (DENV-1 to DENV-4). All four strains leave multiple symptoms including headache, rashes and increased body temperature. Infection and recovery from one strain of the virus can lead to immunity from that particular strain and that issue complicates the modelling of the disease because information on the serotype of infections is rarely available on any wide scale.

Potential individual and ecological risk factors for the disease are varied including both socio-economic and environmental conditions. On the socio-economic front factors such as age, income, population density, sanitation, drainage and water sup-

ply are potentially important. Amongst the environmental considerations, dengue fever is strongly believed to be influenced by climate variability in temperature and precipitation. One useful related measure is also what is commonly referred to as the ‘El Niño Southern Oscillation’ (ENSO) which refers to variations in sea surface temperature (SST) of the tropical eastern Pacific Ocean and in air surface pressure in the tropical western Pacific. ENSO, or equivalently the Oceanic Niño Index (ONI) has three different levels; El Niño, Neutral and La Niña. ONI is a global set of anomalies, and is a useful tool to define patterns of climate change. The most heavy and strong ENSO was reported to occur in the years 1997 to 1998. This El Niño was associated with disasters such as drought, flooding and forest fires around the world (Mark, 2005). Understanding links between ENSO and infectious diseases, particularly those transmitted by insects such as dengue, could provide improved long range forecasting of an epidemic or epizootic (Anyamba et al., 2006). The extent to which ENSO can be linked to epidemics of dengue is still not clear, but there are strong recommendations that it could be investigated in future epidemic forecasting for public health preparedness (Mathuroos et al., 2009).

Studies associating ENSO and other climatic variables to dengue are reviewed in detail in a later chapter of this study; however it is worth making some brief preliminary reference to a selection of some of those here. Studies in Singapore close to Malaysia have been referenced earlier. Farther afield, in Venezuela, Aura and Alfonso (2010) found a significant association between high dengue incidence and lower values of ONI, but lower dengue incidence with higher value of ONI. Meanwhile, in Puerto Rico, Jury (2008) and Earnest et al. (2012b) concluded that the variability of dengue cases was positively related to temperature but weakly associated with local rainfall and ENSO. Hurtado-Diaz et al. (2007) reported every degree increase in SST leading to a 46% increase in dengue cases in San Andre’s Tuxtla and 42% in Veracruz for 16 and 20 weeks respectively. Adriana et al. (2012) used Poisson and Negative Binomial GLMs to investigate the effect of seasonal factors and the relationship of climatic variables to dengue counts in Rio de Janeiro in Brazil. The results indicated significant relationships with the minimum

temperature and precipitation at lag one month before, with a 1 °C increase in a month's minimum temperature leading to a 45% increased in dengue cases in the following month and a 10-millimeter increase in precipitation leading to a 6% increase in dengue in the following month. Other research in the South East of Brazil (Lowe et al., 2013) used Negative Binomial generalised linear mixed models (GLMM) to relate monthly dengue incidence to climate and non-climate covariates. Their results provided probabilistic predictions of future epidemics of dengue several months ahead and the general modelling framework used could apply to other areas of Brazil and other climate sensitive diseases.

One issue which these studies perhaps emphasise is that whilst there is broad agreement that climatic factors do influence variability in dengue incidence, there is no clear consensus as to the degree of such effects or indeed, in some cases, their direction. How climate contributes to increase or decrease the incidence of vector borne diseases in human populations will depend on local climatic conditions and local non-climatic epidemiologic and ecologic factors (Patz and Olson, 2006). In other words, effects are geographically dependent upon the region of the world in question and are confounded with other non-climatic influences in ways which are possibly also geographically specific. It is clear therefore that one cannot necessarily transfer results from elsewhere in the world directly into the Malaysian context. Rather there is a requirement to explore from scratch climatic and other relationships with dengue in the specific Malaysian context if progress is to be made towards developing dengue EWS in Malaysia.

Turning briefly to that Malaysian context (a topic which is picked up in more detail in a subsequent chapter), there are factors which have been suggested globally as encouraging dengue spread which particularly pertain in that country, such as rapid and relatively unorganised urbanisation and high rates of population growth. The rise in global commerce and tourism, global warming and changes in public health policy could be important factors too (Gubler, 1998). Developing economies such as Malaysia are also often criticised for poor construction planning which then causes floods or droughts through failure to consider climate information adequately.



Studies to assess the level of knowledge, attitude and practices in relation to dengue in Malaysia have been conducted in 2003 and 2006 by Hairi et al. (2003) and Wan Rozita et al. (2006). In such studies the aims are to evaluate dengue control through increasing the health promotion activities and exposure of communities to educational campaigns. Results obtained were mixed but generally not very encouraging. Meanwhile, Shekhar and Huat (1992) have highlighted major weaknesses of current epidemiological research on dengue in Malaysia which include the inadequacy of data and lack of sound statistical methods. They considered available data used so far to be too restricted, collected using methods that are not clearly described, and which lack scientific validity. The public health sector at the international level has recognised geographic information systems (GIS)<sup>1</sup> as a new technology which has an ability to change the health of societies and contribute to public health policy investigation, development and execution. The WHO has reported that GIS are potentially valuable tools in data compilation and presentation especially for environmental data linked to health services. In Malaysia, this has been explored in relation to dengue by Shaharudin et al. (2002) with results showing no significant difference in the geographical distribution of dengue cases between 1999 and 2000.

As regards the few studies more directly relevant to this study, Lam (1993b) believed that it is possible to predict the severity of a dengue epidemic by the strain of the circulating serotype, but Chew et al. (2012) makes it clear that the situation in Malaysia is complex – although his study showed the predominance of dengue virus in the capital city of Malaysia (Kuala Lumpur) was DENV-4, it was also the case that all four dengue serotypes were in circulation. Ibrahim et al. (2011) used five years data of dengue (2007-2011) to simulate a dynamic system to predict the spread of dengue outbreak in Hulu Langat, Selangor, Malaysia and the results showed that mean temperature, total amount of rainfall and the total of dengue cases in the previous period were highly significant in predicting the possibility of a dengue outbreak.

---

<sup>1</sup><http://www.gis.com/>



## References

- Abu Bakar, A., Kefli, Z., Abdullah, S., and Sahani, M. (2011). Predictive models for dengue outbreak using multiple rulebase classifiers. *International Conference on Electrical Engineering and Informatics (ICEEI)*, 17-19 July 2011, page doi:10.1109/ICEEI.2011.6021830.
- Abu Bakar, A. L. (1996). *Melaka dan Arus Gerakan Kebangsaan Malaysia*. Penerbit Universiti Malaya, Kuala Lumpur.
- Abu Bakar, S. and Shafee, N. (2002). Outlook of dengue in Malaysia: a century later. *The Malaysian Journal of Pathology*, 24(1):23–27.
- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P. P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P., and Nelkin, E. (2003). The Version-2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis (1979-Present). *Journal Hydrometeorology*, 4(6):1147–1167.
- Adriana, F. G., Aline, A. N., and Oswaldo, G. C. (2012). Temporal analysis of the relationship between dengue and meteorological variables in the city of Rio de Janeiro, Brazil, 2001-2009. *Cad. Saude Publica, Rio de Janeiro*, 28(11):2189–2197.
- Ahmed, S. A., Siddiqi, J. S., Quaiser, S., and Kamal, S. (2015). Using PCA, Poisson and Negative Binomial model to study the climatic factor and dengue fever outbreak in Lahore. *Journal of Basic and Applied Sciences*, 11:8–16.

- Aiken, S. R., Frost, D. B., and Leigh, C. H. (1980). Dengue hemorrhagic fever and rainfall in Peninsular Malaysia: some suggested relationships. *Social Science and Medicine*, 14d(3):307–316.
- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. *Proceeding of the Second International Symposium on Information Theory*, pages 199–213.
- Althouse, B. M., Ng, Y. Y., and Cummings, D. A. T. (2011). Prediction of dengue incidence using search query surveillance. *PLoS Neglected Tropical Diseases*, 5(8):doi:10.1371/journal.pntd.0001258.
- Ang, K. T., Rohani, I., and Look, C. H. (2010). Role of primary care providers in dengue prevention and control in the community. *Medical Journal Malaysia*, 65(1):58–62.
- Anyamba, A., Chretien, J., Small, J., Tucker, C. J., and Linthicum, K. J. (2006). Developing global climate anomalies suggest potential disease risks for 2006/2007. *International Journal of Health Geographics*, 60(5).
- Ashford, D. A., Savage, H. M., Hajjeh, R. A., Mcready, J., Bartholomew, D. M., Spiegel, R. A., Vorndam, V., Clark, G. G., and Gubler, D. G. (2003). Outbreak of dengue fever in Palau, Western Pacific: Risk factors for infection. *American Journal Tropical Medicine Hygiene*, 69(2):135–140.
- Aura, D. H. M. and Alfonso, J. R. M. (2010). Potential influence of climate variability on dengue incidence registered in a western pediatric Hospital of Venezuela. *Tropical Biomedicine*, 27(2):280–286.
- Aziz, A. A. A., Ibrahim, H., and Abdullah, R. (2011). A model of knowledge mapping in visualizing the hotspot of dengue. *IEEE Conference on Open Systems(ICOS2011)*, pages 444–449.
- Aziz, S., Aidil, R. M., Nisfarizal, M. N., Ngui, R., Lim, Y. A. L., Wan Yusoff, W. S., and Wan Ruslan, R. (2014). Spatial density of Aedes distribution in urban

- areas: A case study of breteau index in Kuala Lumpur, Malaysia. *Journal of Vector Borne Disease*, 51:91–96.
- Aziz, S., Ngui, R., Lim, Y. A. L., Solehah, I., Nur Farhana, J., Azizan, A. S., and Wan Yusoff, W. S. (2012). Spatial pattern of 2009 dengue distribution in Kuala Lumpur using GIS application. *Tropical Biomedicine*, 29(1):113–120.
- Azmawati, M. N., Aniza, I., and Ali, M. (2013). Evaluation of Communication for Behavioral Impact (COMBI) Program in Dengue Prevention : A Qualitative and Quantitative Study in Selangor, Malaysia. *Iranian Journal Public Health*, 42(5):538–539.
- Breslow, N. E. (1984). Extra-Poisson variation in log-linear models. *Journal of the Royal Statistical Society*, 33(1):38–44.
- Breslow, N. E. (1996). Generalized linear models checking assumptions and strengthening conclusions. *Congresso Nazionale Societa Italiana di Biometria*, pages 1–18.
- Cabrera, M. (2013). *Spatio-temporal modelling of dengue fever in Zulia state, Venezuela*. PhD thesis, University of Bath, Department of Health, Bath, United Kingdom.
- Carbajo, A. E., Schweigmann, N., Curto, S. I., de Garin, A., and Bejaran, R. (2001). Dengue transmission risk maps of Argentina. *Tropical Medicine and International Health*, 6(3):170–183.
- Chambers, J. M. and Hastie, T. J. (1992). *Statistical Models in S*. Chapman and Hall, London.
- Chan, Y. C., Ho, B. C., and Chan, K. L. (1971). *Aedes aegypti*(L.) and *Aedes albopictus*(Skuse) in Singapore City. *Bulletin World Health Organization*, 44:651–657.
- Chee, H. Y. and Abu Bakar, S. (2003). Phylogenetic investigation of dengue virus type 2 isolated in Malaysia. *Dengue Bulletin*, 27:100–107.

- Chen, C. C. and Chang, H. C. (2013). Predicting dengue outbreaks using approximate entropy algorithm and pattern recognition. *Journal of Infection*, pages 1–7.
- Chen, C. D., Nazni, W. A., Lee, H. L. Seleena, B., Mohd Masri, S., Chiang, Y. F., and Sofian Azirun, M. (2006). Mixed breeding of *Aedes aegypti*(L.) and *Aedes albopictus* skuse in four dengue endemic areas in Kuala Lumpur and Selangor, Malaysia. *Tropical Biomedicine*, 23(2):224–227.
- Cheong, Y. L., Burkart, K., Leitao, P. J., and Lakes, T. (2013). Assessing Weather Effects on Dengue Disease in Malaysia. *International Journal of Environmental Research and Public Health*, 10(12):6319–6334.
- Chew, M. H., Rahman, M. M., Jelip, J., Hassan, M. R., and Isahak, I. (2012). All serotypes of dengue viruses circulating in Kuala Lumpur, Malaysia. *Current Research Journal of Biological Sciences*, 4(2):229–234.
- Choy, E. A., Mohd Khair, E., Atan, A., Sahani, M., and Mohd Ali, Z. (2011). Climate change and dengue: Case study in Seremban District, Negeri Sembilan, Malaysia. *Jurnal e-Bangi*, 6(1):38–48.
- Chuang, P. H., Chuang, J. H., and Lin, I. F. (2010). A dynamic estimation of the daily cumulative cases during infectious disease surveillance: Application to dengue fever. *BMC Infectious Diseases*, 10(136):doi: 10.1186/1471-2334-10-136.
- Cleveland, W. S. and Devlin, S. J. (1998). Locally weighted regression: An approach to regression analysis by local fitting. *Journal of the American Statistical Association*, 83(403):596–610.
- Cleveland, W. S., Grosse, E., and Shyu, W. M. (1992). *Local regression models. Chapter 8 of Statistical Models in S*. Wadsworth and Brooks-Cole.
- Connor, S. J., Omumbo, J., Green, C., DaSilva, J., Mantilla, G., Delacollette, C., Hales, S., Rogers, D., and Thomson, M. (2010). Health and Climate Needs. *Procedia Environmental Sciences*, 1(3):27–36.

- Costello, A., Abbas, M., Allen, A., Ball, S., Bell, S., Bellamy, R., Friel, S., Groce, N., Johnson, A., Kett, M., Lee, M., Levy, C., Maslin, M., McCoy, D., McGuire, B., Montgomery, H., Napier, D., Pagel, C., Patel, J., Antonio, J., Oliveira, P., Redclift, N., Rees, H., Rogger, D., Scott, J., Stephenson, J., Twigg, J., Wolff, J., and Patterson, C. (2009). Managing the health effects of climate change. *www.thelancet.com*, 373:1693–1733.
- Craven and Wahba (1979). Smoothing noisy data with spline functions. *Numerical Mathematics*, 31:377–403.
- Cuadras, C. M. and Fortiana, J. (2002). *Distributions with given marginals and statistical modelling*. Springer.
- Degallier, N., Favier, C., Menkes, C., Lengaigne, M., Ramalho, W. M., Souza, R., Servain, J., and Boulanger, J. P. (2010). Toward an early warning system for dengue prevention: Modeling climate impact on dengue transmission. *Climatic Change*, 98:581–592.
- Dhang, C. C., Benjamin, S., Saranum, M. M., Fook, C. Y., Lim, L. H., Ahmad, N. W., and Azirun, M. S. (2005). Dengue vector surveillance in urban residential and settlement areas in Selangor, Malaysia. *Tropical Biomedicine*, 22(1):39–43.
- Donald, S. S., Rosemary, L., Chiu, W. N., Eduardo, A. U., Yara, H., and Lucy, L. ad Laurent, C. (2012). Burden of dengue in Malaysia. Report from a Collaboration between Universities and the Ministry of Health Malaysia. *Brandeis University*, 42:1–78.
- Earnest, A., Tan, S. B., Wilder-Smith, A., and Machin, D. (2012a). Comparing Statistical Models to Predict Dengue Fever Notifications. *Computational and Mathematical Methods in Medicine*, 2012:doi.org/10.1155/2012/758674.
- Earnest, A., Tana, S. B., and Wilder Smith, A. (2012b). Meteorological factors and El Niño Southern Oscillation are independently associated with dengue infections. *Epidemiology and Infection*, 140(7):1244–1251.

- Ebi, K. L., Hartman, J., Chan, N., McConnell, J., Schlesinger, M., and Weyant, J. (2005). Climate suitability for stable malaria transmission in Zimbabwe under different climate change scenarios. *Climate Change*, 73:375–393.
- Economy, U. (2010). MDG1: Eradicate extreme poverty and hunger. *Malaysia: The Millennium Development Goals at 2010*, pages 10–30.
- Emilie, A., Laurent, G., Beat, S., Francois, C., and Louis, L. (2011). Urbanisation and infectious diseases in a globalised world. *www.thelancet.com/infection*, 11.
- Favier, C., Degallier, N., Rosa-Freitas, M. G., Boulanger, J. P., Costa Lima, J. R., Luitgards Moura, J. F., Menke's, C. E., Mondet, B., Oliveira, C., Weimann, E. T. S., and Tsouris, P. (2006). Early determination of the reproductive number for vector-borne diseases: The case of dengue in Brazil. *Tropical Medicine and International Health*, 11(3):332–340.
- Fredolin, T. T. and Liew, J. (2004). Mechanisms of Malaysian rainfall anomalies. *Journal of Climate*, 17:3616–3622.
- Gage, K. L., Burkot, T. R., Eisen, R. J., and Hayes, E. B. (2008). Climate and vectorborne disease. *American Journal of Preventive Medicine*, 35(5):436–450.
- Gagnon, A. S., Bush, A. B. G., and Smoyer Tomic, K. E. (2001). Dengue epidemics and the El Niño Southern Oscillation. *Climate Research*, 19(1):35–43.
- George, R. (1992). Current status of the knowledge of Dengue/DHF/DSS in Malaysia: Clinical Aspects. *Microbiology and Infectious Diseases*, pages 1–5.
- Gharbil, M., Quenel, P., Gustave, J., Cassadou, S., Ruche, G., Girdary, L., and Marrama, L. (2011). Time series analysis of dengue incidence in Guadeloupe, French West Indies: Forecasting models using climate variables as predictors. *BMC Infectious Diseases*, 11(166).

- Ghebreyesus, T. A., Tadesse, Z., Jima, D., Bekele, E., Mihretie, A., Yihdego, Y. Y., Dinku, T., Connor, S. J., and Rogers, D. P. (2009). Using climate information in the health sector. *Field Actions Science Reports*, 2:63–67.
- Glantz, M. H. (2001). *Currents of change: Impacts of El Niño and La Niña on climate and society*. Cambridge University Press.
- Gubler, D. J. (1998). Dengue and dengue hemorrhagic fever. *Clinical Microbiology Reviews*, 11:480–496.
- Gubler, D. J. (2002). Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends in Microbiology*, 10(2):100–103.
- Gubler, D. J. and Rosen, L. (1976). A simple technique for demonstrating transmission of dengue virus by mosquitoes without the use of vertebrate hosts. *The American Journal of Tropical Medicine and Hygiene*, 25(1):146–150.
- Guzman, M. G. and Kouri, G. (2002). Dengue: An update. *The Lancet, Infectious Diseases*, 2(1):33–42.
- Hafiz, H., Shamarina, S., and Nor, R. H. (2012). Risk mapping of dengue in Selangor and Kuala Lumpur, Malaysia. *Geospatial Health*, 7(1):21–25.
- Hairi, F., Ong, C. H. S., Suhaimi, A., Tsung, T. W., Anis Ahmad, M. A., Sundaraj, C., and M., S. M. (2003). A knowledge, attitude and practices (KAP) study on dengue among selected rural communities in the Kuala Kangsar district. *Asia Pacific Journal Public Health*, 15(1):37–43.
- Hales, S., Wet, N. D., Maindonald, J., and Woodward, A. (2002). Potential effect of population and climate changes on global distribution of dengue fever: An empirical model. *The Lancet*, pages 1–5.
- Hashizume, M., Dewan, A. M., Sunahara, T., Rahman, M. Z., and Yamamoto, T. (2012). Hydroclimatological variability and dengue transmission in Dhaka, Bangladesh: a time-series study. *BMC Infectious Diseases*, 12(98):1–9.



- Hastie, T. and Tibshirani, R. (1986). Generalized Additive Models. *Statistical Science*, 1(3):297–318.
- Hastie, T. J. and Tibshirani, R. J. (1990). *Generalized Additive Models*. Chapman and Hall.
- Heil, A. and Goldammer, J. G. (2001). Smoke-haze pollution: A review of the 1997 episode in Southeast Asia. *Regional Environmental Change*, 2(1):24–37.
- Hii, Y. L., Zhu, H., Ng, N., Ng, L. C., and Rocklov, J. (2012). Forecast of dengue incidence using temperature and rainfall. *PLoS Negl Trop Dis*, 6(11):doi: 10.1371/journal.pntd.0001908.
- Hirschman, C. (1987). The meaning and measurement of ethnicity in Malaysia: An analysis of census classifications. *The Journal of Asian Studies*, 46(3):555–582.
- Hopp, J. M. and Foley, J. A. (2003). Worldwide fluctuations in dengue fever case related to climate variability. *Climate Research*, 25(1):85–94.
- Hu, W., Clements, A., William, G., Tong, S., and Mengersen, K. (2012). Spatial patterns and socioecological drivers of dengue fever transmission in Queensland, Australia. *Environmental Health Perspectives*, 120(2):260–266.
- Hurtado-Diaz, M., Riojas-Rodriguez, H., Rothenberg, S. J., Gomez-Dantes, H., and Cifuentes, E. (2007). Short Communication: Impact of climate variability on the incidence of dengue in Mexico. *Tropical Medicine and International Health*, 12(11):1327–1337.
- Ibrahim, A., Mat Zin, N. A., and Sahari Ashaari, N. (2011). Simulation model for predicting dengue fever outbreak. *World Academy of Science, Engineering and Technology*, 5(11).
- Ibrahim, F., Wan Abas, W. A. B., Chong Guan, C., and Sulaiman, S. (2007). A new approach to classify risk in dengue infection using bioelectrical impedance analysis. *Dengue Bulletin*, 31:58–74.



- IPCC (2014). Fifth Assessment Report - Climate Change 2014 : Impacts, Adaptation, and Vulnerability. *Intergovernmental Panel Climate Change*.
- Jaafar, S., Mohd Noh, K., Abdul Mutalib, K., Othman, N. H., and Healy, J. (2013). Malaysia Health System Review. *Health Systems in Transition*, 3(1):82–92.
- Johansson, M. A., Cummings, D. A. T., and Glass, G. E. (2009a). Multiyear climate variability and dengue El Niño Southern Oscillation, weather and dengue incidence in Puerto Rico, Mexico and Thailand: A longitudinal data analysis. *PLoS Medicine*, 6(11):doi: 10.1371/journal.pmed.1000168.
- Johansson, M. A., Dominici, F., and Glass, G. E. (2009b). Local and global effects of climate on dengue transmission in Puerto Rico. *PLoS Neglected Tropical Diseases*, 3(2):doi: 10.1371/journal.pntd.0000382.
- Juneng, K. and Tanggang, F. T. (2008). Level and source of predictability of seasonal rainfall anomalies in Malaysia using canonical correlation analysis. *International Journal of Climatology*, 28:1255–1267.
- Juni, M. H., Hayati, K. S., Cheng, C. M., Pyang, G. S., Nur Hayati, A. S., and Zaidatul Syeila, Z. A. (2015). Risk behaviour associated with dengue fever among rural population in Malaysia. *International Journal of Public Health and Clinical Sciences*, 2(1):114–127.
- Jury, M. R. (2008). Climate influence on dengue epidemics in Puerto Rico. *International Journal of Environmental Health Research*, 18(5):323–334.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Roy, J., and Dennis, J. (1996). The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American Meteorological Society*, 77(3):437–471.

- Khormi, H. M. and Kumar, L. (2011). Modelling dengue fever risk based on socioeconomic parameters, nationality and age groups: GIS and remote sensing based case study. *Science of Total Environment*, 409(22):4713–4719.
- Khormi, H. M., Kumar, L., and Elzahrany, R. A. (2011). Modeling spatio-temporal risk changes in the incidence of dengue fever in Saudi Arabia: A geographical information system case study. *Geospatial Health*, 6(1):77–84.
- Khun, K., Campbell Lendrum, D., Haines, A., Cox, J., Corvalan, C., and Anker, M. (2005). Using climate to predict infectious disease epidemic. *World Health Organization*.
- Kovats, R. S., Bouma, M. J., and Haines, A. (1999). El Niño and Health. *Task Force on Climate and Health, World Health Organization, Geneva*, pages 1–46.
- Krisada, L. and Lily, I. (2013). Generalized linear mixed models with spatial random effects for spatio-temporal data: An application to dengue fever mapping. *Journal of Mathematics and Statistics*, 9(2):137–143.
- Kumarasamy, V. (2006). Dengue fever in Malaysia: Time for review? *Medical Journal Malaysia*, 61(1):1–3.
- Kusriastuti, R. and Sutomo, S. (2005). Evolution of Dengue Prevention and Control Programme in Indonesia. *Dengue Bulletin*, 29:1–7.
- Lam, S. K. (1993a). Strategies for dengue control in Malaysia. *Tropical Medicine*, 35(4):303–307.
- Lam, S. K. (1993b). Two decades of dengue in Malaysia. *Tropical Medicine*, 35(4):195–200.
- Lee, H. L., Rohani, A., Khadri, M. S., Nazni, W. A., Rozilawati, H., Nurulhusna, A. H., Nor Afizah, A. H., Roziyah, A., Rosilawati, R., and Teh, C. H. (2015). Dengue Vector Control in Malaysia-Challenges and Recent Advances. *The International Medical Journal Malaysia*, 14(1):11–16.

- Loevinsohn, M. E. (1994). Climatic warming and increased malaria incidence in Rwanda. *The LANCET*, 343:714–718.
- Loh, B. and Song, R. J. (2001). Modeling dengue cluster size as a function of *Aedes aegypti* population and climate in Singapore. *Dengue Bulletin*, 25:74–78.
- Lowe, R. (2010). *Spatio-temporal modelling of climate-sensitive disease risk: Towards an early warning system for dengue in Brazil*. PhD thesis, College of Engineering, Mathematics and Physical Science, University of Exeter, Harrison Building, College of Engineering, Mathematics and Physical Science, EX4 4QF, Exeter, United Kingdom.
- Lowe, R., Bailey, T. C., Stephenson, D. B., Jupp, T. E., Graham, R. J., Barcellos, C., and Carvalho, M. S. (2013). The development of an early warning system for climate-sensitive disease risk with a focus on dengue epidemics in Southeast Brazil. *Statistics in Medicine*, 32(5):864–883.
- Lu, L., Lin, H., Tian, L., Yang, W., Sun, J., and Liu, Q. (2009). Time series analysis of dengue fever and weather in Guangzhou, China. *BMC Public Health*, 9:395–399.
- Ma, S., Ooi, E. E., and Goh, K. T. (2008). Socioeconomic determinants of dengue incidence in Singapore. *Dengue Bulletin*, 32.
- Mammen, M. P. J., Pingate, C., Koenraadt, C. J. M., Rothman, A. L., Aldstadt, J., Nisalak, A., Jarman, R. G., Jones, J. W., Srikiatkachorn, A., Ypil Butac, C. A., Getis, A., Thammapalo, S., Morrison, A. C., Libraty, D. H., Green, S., and Scott, T. W. (2008). Spatial and temporal clustering of dengue virus transmission in Thai Villages. *PLoS Medicine*, 5(11):1605–1615.
- Mark, A. C. (2005). The evolution of El Niño, past and future. *Earth and Planetary Science Letters*, 230:227–240.
- Markon, A. O. (2014). *Dengue fever in Ribeirao Preto, Brazil, 2003-2012: Patterns of Disease and Understanding of Prevention*. PhD thesis, University of Michigan, University of Michigan, United States.

- Martens, W. J. M., Niessen, L. W., Rothmans, J., Jetten, T. H., and McMichael, A. J. (1995). Potential impact of global climate change on malaria risk. *Environmental Health Perspectives*, 103(5):458–464.
- Mary Ann, B. (2009). Climate change could affect mosquito-borne diseases in Asia. *www.thelancet.com*, 373:1070.
- Mathuros, T., Chi Tai, F., Suratsawadee, K., Chung Ming, L., and Chwan Chuen, K. (2009). Effects of the El Niño-Southern Oscillation on dengue epidemics in Thailand, 1996-2005. *BMC Public Health*, 9:422–436.
- Mazrura, S., Rozita, H., Hidayatulfathi, O., Zainudin, M. A., Mohamad Naim, M. R., Nadia Atiqah, M. N., Rafeah, M. N., Er, A., Norela, S., Nurul Ashikin, Z., and Joy, J. P. (2010). Community Vulnerability On Dengue And Its Association With Climate Variability in Malaysia: A Public Health Approach. *Malaysian Journals of Public Health Medicine*, 10(2):25–34.
- McCullagh, P. and Nelder, J. A. (1989). Generalized linear models. *Ecology*, 88(11).
- Mia, M. S., Begum, R. A., Er, A. C., Abidin, R. D., and Pereira, J. J. (2013). Trends of dengue infections in Malaysia, 2000-2010. *Asian Pacific Journal of Tropical Medicine*, 6(6):462–466.
- Ministry, H. M. (2008). Health Facts 2008 Malaysia. *Ministry of Health Malaysia*, pages 1–12.
- Ministry, H. M. (2014). Health Facts 2014 Malaysia. *Ministry of Health Malaysia*, pages 1–12.
- Mohamad Ismail, N. A., Kampan, N., Abdullah Mahdy, Z., Abdul Jamil, M., and Mohd Razi, Z. R. (2011). Dengue in pregnancy. *Southeast Asian Journal Trop Med Public Health*, 37(4):681–683.
- Mohamad Mohsin, M. F., Abu Bakar, A., and Hamdan, A. R. (2013). The Preliminary Design of Outbreak Detection Model Based on Inspired Immune

- System. *Third World Congress on Information and Communication Technologies (WICT)*, pages 189–195.
- Mohd Ismail, M. (2007). Malaysia's Health 2007. *Kementerian Kesihatan Malaysia*, pages 1–299.
- Mohd Ismail, M., Ahmad, K., Kamarul Zaman, M. I., Ramlee, R., Noor Hisham, A., Maimunah, A. H., Norain, A. T., Che Mohd Zain, C. A., Abd Rahim, M., and Khadzir, S. A. (2007). Laporan Tahunan 2007. *Kementerian Kesihatan Malaysia*, pages 1–293.
- Mohd Ismail, M., Ahmad Shafii, S., Kamarul Zaman, M. I., Ramlee, R., Noor Hisham, A., Maimunah, A. H., Norain, A. T., Eisah, A. R., Abd Rahim, M., and Khadzir, S. A. (2009). Laporan Tahunan 2009. *Kementerian Kesihatan Malaysia*, pages 1–345.
- Monath, T. P. (1994). Dengue: the risk to developed and developing countries. *Proceedings of the National Academy of Science of the United States of America*, 91(7):2395–2400.
- Mondini, A. and Chiaravalloti, N. F. (2007). Socioeconomic variables and dengue transmission. *Rev Saude Publica*, 41(6):923–930.
- Muhamed, K. and Haron, S. (2011). Poverty Mapping: An Effective Approach in Determining The Poor Area - Case Study of Johor. *Department of Statistics, Malaysia*, pages 43–60.
- Muhammad Azami, N. A., Salleh, S. A., Neoh, H. M., Syed Zakaria, S. Z., and Jamal, R. (2011). Dengue epidemic in Malaysia: not a predominantly urban disease anymore. *BMC Research Notes*, 216(4):1–4.
- Narwani, H., Jesni, J., Nyi Nyi, N., Hamzah, A. M., Abd Haris, M., and Mohd Noor, M. (2005). A review of dengue fever incidence in Kota Bharu, Kelantan, Malaysia during the years 1998–2003. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 36(5):1179–1186.

- Natchaporn, P., Noparat, M., Siripen, K., and Wisit, C. (2006). Relationship between body size and severity of dengue hemorrhagic fever among children aged 0-14 years. *Southeast Asian Journal Tropical Medical Public Health*, 37(2):283–288.
- Nelder, J. A. and Wedderburn, R. W. M. (1972). Generalized linear models. *Journal of the Royal Statistical Society*, 135(3):370–384.
- Nicol, J. (1997). Bioclimatic impacts of the 1994 smoke-haze event in Southeast Asia. *Atmospheric Environment*, 31(8):1209–1219.
- Nor Azimi, Y. (2000). Dengue outbreak in 1998-Terengganu experience. *Buletin Kesihatan Masyarakat Isu Khas*, pages 75–84.
- Oki, T. and Musiake, K. (1994). Seasonal change of the diurnal cycle of precipitation over Japan and Malaysia. *American Meteorological Society*, 33:1445–1463.
- Ooi, E. E. and Gubler, D. J. (2008). Dengue in Southeast Asia: epidemiological characteristics and strategic challenges in disease prevention. *Cadernos de Saude Publica*, 25:115–124.
- Osgood, D. W. (2000). Poisson-based regression analysis of aggregate crime rates. *Journal of Quantitative Criminology*, 16(1):21–43.
- Osman, O., Fong, M. Y., and Devi, S. (2007). A preliminary study of dengue infection in Brunei. *Journal Infectious Diseases*, 60:205–208.
- Parks, W. J., Lloyd, L. S., Nathan, M. B., Hosein, E., Odugleh, A., Clark, G. G., Gubler, D. J., Palmer, K., Martin, J. L. S., Siversen, S. R., Dawkins, Z., and Renganathan, E. (2004). International experiences in social mobilization and communication for dengue prevention and control. *Dengue Bulletin*, 28:1–7.
- Patz, J. A., Martens, W. J., Focks, D. A., and Jetten, T. H. (1998). Dengue fever epidemic potential as projected by general circulation models of global climate change. *Environmental Health Perspectives*, 106(3):147–153.



- Patz, J. A. and Olson, S. H. (2006). Climate change and health: Global to local influences on disease risk. *Annals of Tropical Medicine and Parasitology*, 100(5):535–549.
- Poovaneswari, S. (1993). Dengue situation in Malaysia. *Malaysian Journal Pathology*, 15(1):3–7.
- Pouliot, S. H., Xiong, X., Harville, E., Paz Soldan, V., Tomashek, K., Breart, G., and Buekens, P. (2010). Maternal dengue and pregnancy outcomes: A systematic review. *Obstetrical and Gynecological Survey*, 65(2):69–144.
- R Core Team, D. (2010). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Racloz, V., Ramsey, R., Tong, S., and Hu, W. (2012). Surveillance of dengue fever virus: A review of epidemiological models and early warning systems. *PLoS Neglected Tropical Diseases*, 6(5):doi: 10.1371/journal.pntd.0001648.
- Richard, B. and John, M. (2007). Overdispersion and Poisson Regression. *Social and Economic Sciences*, pages 1–24.
- Rogers, D. P., Boulahya, M. S., Thompson, M. A., Connor, S. J., Dinku, T., Johm, K. B., Shalaby, H. R., Ahmadu, B., and Niang, A. (2008). National climate and environmental service for development: Proceedings of the International Symposium on PWS: A Key to Service Delivery. *World Meteorological Organization*, pages 87–92.
- Rosnah, H., Za'aba, B., Abdul Rasid, K., Maimunah, A. R., Yogeswary, M., Mohd Jubri, J., Noorazah, D., Khairul Rafizah, H., Siti Yuhanis, I., Rafidah, S., Norhayati, M. S., Mas'ud, H., Rasyikah, M., Rajini, M., and Tg Mawar, T. A. A. (2009). Laporan Tahunan 2009. *Jabatan Kesihatan Negeri Selangor*, pages 1–180.
- Rowley, W. A. and Graham, C. L. (1968). The effect of temperature and relative humidity on the flight performance of female *Aedes aegypti*. *Journal Insect Physiol.*, 14:1251–1257.

- Rozhan, S., Jamsiah, M., Rahimah, A., and Ang, K. T. (2006). The COMBI (Communication for Behavioral Impact) Program in the prevention and control of dengue-The Hulu Langat experience. *Jurnal Kesehatan Masyarakat*, 12(1):19–32.
- Rozilawati, H., Zairi, J., and Adanan, C. R. (2007). Seasonal abundance of *Aedes albopictus* in selected urban and suburban areas in Penang, Malaysia. *Tropical Biomedicine*, 24(1):83–94.
- Rozita, H., Hidayatulfathi, O., Nurul Azimah, J., Mazrura, S., Kamarulismail, U., Zainudin, M. A., Er, A. C., and Zailiza, S. (2013). The COMBI Approach in Managing Dengue Cases in an Urban Residential Area, Nilai, Malaysia. *International Journal of Public Health Research*, 3(2):347–352.
- Rudnick, A., Tan, E. E., Lucas, J. K., and Omar, M. (1965). Mosquito-borne haemorrhagic fever in Malaya. *British Medical Journal*, 1(5445):1269–1272.
- Russell, R. C., Currie, B. J., Lindsay, M. D., Mackenzie, J. S., Ritchie, S. A., and Whelan, P. I. (2009). Dengue and climate change in Australia: Predictions for the future should incorporate knowledge from the past. *The Medical Journal of Australia*, 190(5):265–268.
- Sakamoto, Y., Ishiguro, M., and Kitagawa, G. (1988). *Akaike Information Criterion Statistics*. KTK Scientific Publisher.
- Schreiber, K. V. (2001). An investigation of relationships between climate and dengue using a water budgeting technique. *International Journal of Biometeorology*, 45(2):81–89.
- Schwartz, E. (2002). Study of dengue fever among Israeli travellers to Thailand. *Dengue Bulletin*, 26:162–167.
- Schwartz, G. (1978). Estimating the dimension of a model. *Annals of Statistics*, 6:461–464.



- Seng, S. B., Chong, A. K., and Moore, A. (2005). Geostatistical modelling, analysis and mapping of epidemiology of dengue fever in Johor State, Malaysia. *The 17th Annual Colloquium of the Spatial Information Research Centre*, pages 1–14.
- Setyono, A., Dian, N., Alam, M. J., and Soosai, B. A. M. (2011). Study and design framework for e-Dengue teleconsultation system. *Instrumentation, Communications, Information Technology, and Biomedical Engineering (ICICI-BME)*, pages 293–297.
- Shaharudin, I., Shamsul, A. S., Tahir, A., Mariam, M., Azah, D., and Nik Shamsidah, N. I. (2002). Sistem maklumat geografi (GIS) dan sektor kesihatan awam: Kajian demam denggi di Bandar Baru Bangi dan Kajang. *Jurnal Kesihatan Masyarakat Isu Khas*.
- Shekhar, K. C. and Huat, O. L. (1992). Epidemiology of dengue/dengue hemorrhagic fever in Malaysia-A retrospective epidemiological study 1973-1987. Part I: Dengue hemorrhagic fever (DHF). *Asia Pacific Journal of Public Health*, 6(2):15–25.
- Simões, T. C., Codeco, C. T., Nobre, A. A., and Eiras, A. E. (2013). Modeling the non-stationary climate dependent temporal dynamics of *Aedes Aegypti*. *PLoS ONE*, 8(8):1–10.
- Simon, P. W., Matthew, G. K., and Fred, L. M. (2003). *Statistical and econometric methods for transportation data analysis*. Chapman and Hall/Crc.
- Simon, W. (2006). *Generalized Additive Models: An introduction with R*. Chapman and Hall.
- Singh, N. P., Jhamb, R., Agarwal, S. K., Gaiha, M., Dewan, R., Daga, M. K., Chakravarti, A., and Kumar, S. (2005). The 2003 outbreak of dengue fever in Delhi India. *Southeast Asian Journal Tropical Medical Public Health*, 36(5):1174–1178.

- Siqueira, J. B., Martelli, C. M. T., Coelho, G. E., Rocha Simplicio, A. C., and Hatch, D. L. (2005). Dengue and dengue hemorrhagic fever, Brazil, 1981–2002. *Emerging Infectious Diseases*, 11(1):48–53.
- Siqueira, J. J. B., Maciel, I. J., Barcellos, C., Souza, W. V., Carvalho, M. S., Nascimento, N. E., Oliveira, R. M., Morais-Neto, O., and Martelli, C. M. T. (2008). Spatial point analysis based on dengue surveys at household level in Central Brazil. *BMC Public Health*, 8:361–369.
- Siwar, C., Ahmed, F., and Mohd Idris, N. D. (2013). Relationship between Poverty and Socio-demographic Characteristics of Poor Households : A Case Study in Kelantan, Malaysia. *2013 International Conference on Advances in Social Science, Humanities and Management (ASSHM 2013)*, 46:679–684.
- Skae, F. M. T. (1902). Dengue fever in Penang. *British Medical Journal*, 2(2185):1581–1583.
- Smith, C. E. G. (1957). A localized outbreak of dengue fever in Kuala Lumpur: Serological aspects. *The Journal of Hygiene*, 55(2):207–223.
- Sokrin, K. and Lenore, M. (2007). Health seeking and access to care for children with suspected dengue in Cambodia: An ethnographics study. *BMC Public Health*, 7:262–271.
- Soman, M. K. and Slingo, J. (1997). Sensitivity of the Asian summer monsoon to aspects of sea surface temperature anomalies in the tropical Pacific ocean. *Quarterly Journal of the Royal Meteorological Society*, 123(538):309–336.
- Souza, S. S., Silva, I. G., and Silva, H. H. (2010). Association between dengue incidence, rainfall and larval density of *Aedes aegypti*, in the State of Goias. *Revista da Sociedade Brasileira Medicina Tropical*, 43(2):152–155.
- Stefan, M., Eng, E. O., and Kee, T. G. (2008). Socioeconomic determinants of dengue incidence in Singapore. *Dengue Bulletin*, 32:17–28.

- Strickman, D. and Kittayapong, P. (2003). Dengue and its vector in Thailand: Calculated transmission risk from total pupae counts of *Aedes aegypti* and association of wing-length measurement with aspects of the larval habitat. *The American Society of Tropical Medicine and Hygiene*, 68(2):209–217.
- Suhaila, J., Deni, S. M., Wan Zin, W. Z., and Jemain, A. A. (2010a). Spatial patterns and trends of daily rainfall regime in Peninsular Malaysia during the southwest and northeast monsoons: 1975-2004. *Meteorology and Atmospheric Physics*, 110(1):doi-10.1007/s00703-010-0108-6.
- Suhaila, J., Deni, S. M., Wan Zin, W. Z., and Jemain, A. A. (2010b). Trends in Peninsular Malaysia rainfall data during the southwest monsoon and northeast monsoon seasons: 1975-2004. *Sains Malaysiana*, 39(4):533–542.
- Suhaili, M. R., Hosein, E., Mokhtar, Z., Ali, N., Palmer, K., and Isa, M. M. (2004). Applying Communication-for-Behavioural-Impact (COMBI) in the prevention and control of dengue in Johor Bahru, Johore, Malaysia. *Dengue Bulletin*, 28:39–43.
- Tanggang, F., Latif, M. T., and Juneng, L. (2010). The roles of climate variability and climate change on smoke haze occurrences in Southeast Asia region . *Climate change: Is Southeast Asia up to the challenge*, 32(5):36–49.
- Tanggang, F. T. and Bahari, A. (2002). ENSO influences on precipitation and air temperature variability in Malaysia. *Proceedings of the Regional Symposium on Environment and Natural Resources*, 1:124–131.
- Teixeira, M. G., Costa, M. C. N., Guerra, Z., and Barreto, M. L. (2006). Dengue in Brazil: Situation-2001 and trends. *Dengue Bulletin*, 26:70–76.
- Teng, A. K. and Singh, S. (2001). Epidemiology and new initiatives in the prevention and control of dengue in Malaysia. *Dengue Bulletin*, 25:7–14.
- Thammapalo, S., Chongsuwatwong, V., Mc Neil, D., and Geater, A. (2005). The climatic factors influencing the occurrence of dengue hemorrhagic fever

- in Thailand. *Southeast Asian Journal Tropical Medicine Public Health*, 36(1):191–196.
- Thomas, W. Y. and Neil, D. M. (1991). Generalized additive models in plant ecology. *Journal of Vegetation Science*, 2(5):587–602.
- Tren, R. and Bate, R. (2001). *Malaria and the DDT Story*. The Institute of Economic Affairs 2001.
- Wahba (1990). *Spline models of observational data*. SIAM.
- Wan Abd Raof, W. R. (2010). *Statistics Handbook Malaysia 2010*. Department of Statistics Malaysia, Pesuruhjaya Banci, Malaysia.
- Wan Abd Raof, W. R. (2011). *Population Distribution and Basic Demographic Characteristics*. Department of Statistics Malaysia, Pesuruhjaya Banci, Malaysia.
- Wan Fairos, W. Y., Wan Azaki, W. H., Mohamad Alias, L., and Bee Wah, Y. (2010). Modelling dengue fever (DF) and dengue haemorrhagic fever (DHF) outbreak using poisson and negative binomial model. *World Academy of Science, Engineering and Technology*, 62:903–908.
- Wan Norafikah, O., Chen, C. D., Soh, H. N., Lee, H. L., Nazni, W. A., and Sofian Azirun, M. (2009). Surveillance of *Aedes* mosquitoes in a university campus in Kuala Lumpur, Malaysia. *Tropical Biomedicine*, 26(2):206–215.
- Wan Rozita, W. M., Yap, B. W., Veronica, S., Muhammad, A. K., Lim, K. H., and Sumarni, M. G. (2006). Knowledge, attitude and practices (KAP) survey on dengue fever in an urban malay residential area in Kuala Lumpur. *Malaysian Journal of Public Health Medicine*, 6(2):62–67.
- Webster, P. J. and Palmer, T. N. (1997). The past and future of El Niño. *Nature*, 390:562–564.

- WHO (2012a). Communication for behavioural impact (COMBI): A toolkit for behavioural and social communication in outbreak response. *Geneva: World Health Organization (WHO)*.
- WHO (2012b). Dengue and severe dengue. *Fact sheet*.
- Wolf, P. S., Motoi, S., Vu, D. T., Richard, G. W., Ataru, T., Lay, M. Y., Hideki, Y., Ubydul, H., Le, H. T., Dang, D. A., and Koya, A. (2011). Population density, water supply, and the risk of dengue fever in Vietnam: Cohort study and spatial analysis. *PLoS Med*, 8(8):1–10.
- Wong, C. L., Venneker, R., Uhlenbrook, S., Jamil, A. B. M., and Zhou, Y. (2009). Variability of rainfall in Peninsular Malaysia. *Hydrol. Earth Syst. Sci.*, 6:5471–5503.
- Wu, P. C., Guo, H. R., Lung, S. C., Lin, C. Y., and Su, H. J. (2007). Weather as an effective predictor for occurrence of dengue fever in Taiwan. *Acta Tropica*, 103(1):50–57.
- Yap, H. H., Chong, N. L., Foo, A. E. S., and Lee, C. Y. (1994). Dengue vector vontrol: Present status and future prospects. *Kaohsiung Journal Medical Science*, 10:S102–S108.
- Yara, A. H., Vishal, D., Narendra, A., Tyagi, B. K., Deoki, N., and Donald, S. S. (2011). Overcoming data limitations: Design of a multicomponent study for estimating the economic burden of dengue in India. *Dengue Bulletin*, 35:1–14.
- Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., and Smith, G. M. (2009). *Mixed Effects Models and Extensions in Ecology with R*. Springer Science, New York.